

Properties and Development of Valortim™ (MDX-1303): A potent, fully human mAb to PA for treatment of inhalational anthrax

**TIBOR KELER, PhD, LAURA VITALE, BS, TOM O'NEIL, BS, DIANN
BLANSET, MS, GEOFFREY NICHOL, MD, ISRAEL LOWY, MD, PhD**

March 29, 2006

**1st NIAID Workshop: Monoclonal Antibody Therapeutics
for Biodefense and Emerging Infectious Diseases**


Medarex/PharmAthene Collaboration

- Valortim™ development began in October 2001
 - Collaborators: Dartmouth Medical School, USAMRIID
 - Identification of a fully human anti-PA mAb using Medarex UltiMAb HuMAb Development System®
 - Two NIAID/DMID grants awarded in September 04
 - Challenge grant – Product Development to FTIH
 - VTAD grant – MoA Studies & Therapeutic Models
- In November 2004, Medarex and PharmAthene entered into a collaborative agreement for advanced development and commercialization of Valortim™

Valortim™: Key Features

- Selected on basis of superior performance in *in vitro* TNA
- Defines a novel neutralizing epitope on PA – NOT attachment
- Mechanism of action similar to protective immune response in vaccinated individuals – FcR dependence on activity
- Potent efficacy in prophylactic and also in therapeutic model of rabbit anthrax aerosol challenge
- Potent efficacy as a prophylactic in the model of monkey aerosol challenge
- Fully human IgGκ monoclonal antibody Medarex Ultimab
 - Produced in CHO cells in defined media
- No human safety issues observed in FTIH – fully enrolled
- Process development complete, Valortim™ ready for full scale production

Antibodies to PA As Antitoxins: Multiple Potential Targets




Potential MAb Activity:

1. Attachment
2. Processing
3. Multimerization
4. Binding of EF/LF
5. Internalization
6. Pore formation

Assays for Lead MAb Selection?


- Affinity of Binding to PA-83/ PA-63
 - Highest affinity screening by Biacore, ELISA, etc...
- Blockade of Binding of PA to toxin receptor or to EF/LF
 - Anthrax Toxin Receptor: eg murine MAb 14B7
 - LF or EF component: eg murine MAb 1G3
- Potency in Toxin Neutralization Assay (TNA)
 - High titer immune sera correlate with TNA activity
 - Standard for vaccine response in clinical trials
- We chose to select HuMAbs with the best activity in TNA as leads, and figure out how they worked later

Valortim™ (MDX-1303): Toxin Neutralization Activity



TNA activity of CDC human reference plasma

AVR801 = 109 $\mu\text{g/ml}$ anti-PA IgG



Valortim: ~ 85x > specific activity
relative to AVR801

AVR801 Reference Standard:
pooled polyclonal antisera from
individuals vaccinated with
approved AVA vaccine

TNA Activity Relative to Other Mabs and to Human Reference Standard

Test Article	EC50(ng/ml)
Valortim™	8
1G3	30
AVR801*	680
14B7	1100

* CDC Human Reference Plasma

CDC Evaluation of Valortim™ in TNA

Origin	ID	N	EC50 Mean (ng/ml)	StdDv	Proprietary EC50 (nM)	CDC EC50 (nM)
Medarex	Valortim™	24	6.39	1.77	0.13nM	0.04nM
Human Reference Standard	AVR414	25	131.48	34.17	Not Applicable	0.88nM


CDC Conclusions

Confirmed Valortim™ of high potency with an EC50 value of 6.39ng/ml (0.04nM)

Performance of Valortim™ has highly reproducible (SD of 1.77)

Valortim(MDX-1303) Defines a Novel Neutralizing Epitope

■ Isotype Control ■ Mu MAb no block ■ Mu MAb + MDX-1303



Mu Mab: 14B7

2D5

1G3

binding epitope : 671-721aa

581-601aa

168-314aa

Blocks receptor binding

Block LF binding




PharmAthene


Little, et al Microbiology 1996,142:707-15

MEDAREX

Receptor binding studies with human monocytes




Valortim (MDX-1303) AND AVR801
do NOT block receptor binding!




% blocking

78%




0 %



In Vivo Efficacy Data

Prophylactic Rabbit Inhalation Model: Dose Titration of Valortim (MDX-1303)



Exposure = 100-300 LD₅₀


Valortim™ Rabbit Inhalation Model

Dose Response


Number of Rabbits	1 Hour Post Exposure	72 or 96 Hours Post Exposure	Survival
10	0 mg/kg	0 mg/kg	0/10
10	1 mg/kg	1 mg/kg	9/10
10	3 mg/kg	3 mg/kg	8/10
10	6 mg/kg	3 mg/kg	9/10
10	12 mg/kg	6 mg/kg	8/10

Lowest Effective Dose Not Yet Identified

Rabbit Inhalation Model: Delayed Treatment with Valortim (MDX-1303)



Exposure ~ 300 LD₅₀



Valortim™ Rabbit Inhalation Model

Delayed Treatment

Number of Rabbits	Antibody	Schedule of Treatments	Dose Level (mg/kg IV)		Survival
			1 st	2 nd	
10	Saline	1 hour post-exposure	0	0	0/10
10	Valortim™	24 hours post- exposure	10	10	8/9 ^a
10	Valortim™	48 hours post-exposure	10	10	3/7 ^b

Target LD50 = 200

^a1 rabbit removed from study unrelated to anthrax or treatment


^b3 rabbits died prior to treatment

Survival to day 14

Therapeutic Models are very challenging because of brittle process



Cynomolgus Monkey Survival After Challenge and IM Injection of Valortim



Valortim™ Monkey Inhalation Model


Number of Monkeys	Treatment	Dose Level 1 Hour Post Exposure	Survival
6	Saline	0 mg/kg IM	0/6
6	Valortim™	1 mg/kg IM	6/6
6	Valortim™	10 mg/kg IM	6/6

- Target LD50 = 200
- Survival to day 90 post-exposure
- Unprotected animals died within 4-7 days


Valortim *in vivo* TNA PK profile

Calculated vs Actual TNA activity in uninfected Cynomolgus monkeys

Group 3: 1mg/kg IM



Group 4: 10mg/kg IM



CDC Human Reference Plasma- AVR801 , 109 $\mu\text{g/ml}$ anti-PA IgG



PharmAthene

MEDAREX

Valortim™ First Time in Human Study

Number of Subjects	Dose	Date of Dosing	Number of SAEs
3	0.3 mg/kg IV	28 Oct 05	0
10	1.0* mg/kg IV	29 Nov 05 30 Nov 05	0
10	3.0 mg/kg IV	05 Jan 06 06 Jan 06	0
10	10.0 mg/kg IV	02 Feb 06 03 Feb 06	0
3	20.0 mg/kg IV	02 Mar 06	0
10	100 * mg IM	08 Mar 06 09 Mar 06	0

*Doses of ≥ 1 mg/kg predicted to be fully protective based on non-human primate studies.

Serum PK, TNA and immunogenicity analyses to commence soon

Valortim - MDX-1303

Anti-PA HuMAb

- Identified by functional screen: potent protection of J744A.1 cells from PA-LT complex mediated death (TNA)
- Potent in vitro neutralizing activity,
 - $ED_{50} = 6-8$ ng/ml, $ED_{80} = 30$ ng/ml ($>$ than known murine MAbs)
 - High affinity to PA83& PA63, $KD = 5.3 \times 10^{-9}$ M
 - Novel epitope, near the receptor binding area but does not block
 - Can rescue macrophages after toxin complex exposure
- Potent in-vivo activity in rabbit AND monkey inhalation model:
 - Protects rabbits at 1 mg/kg (lowest tested to date)
 - Rescues symptomatic rabbits (3/7 survive at 48 hrs admin)
 - Fully protects non-human primates at dose of 1 mg/kg IM
 - After 28 days, 1 mg/kg IM in cyno TNA $>$ CDC reference std
 - Does not block endogenous anti-anthrax immune response
- Well tolerated in normal volunteers at doses likely higher than needed




Mechanism of Action

- Optimal activity utilizes FcR binding


In Vitro Toxin Neutralization:

Valortim Activity is Markedly Enhanced by FcR Interaction




FcR Augmentation of Toxin Neutralization: Found in Some, But Not All MAbs

Blocking of Toxin Neutralization




Toxin Neutralization by Polyclonal Anti-PA Sera: Markedly Augmented by FcR Interaction

Mouse




Rabbit



Anti-Fc receptor mAb 2.4G2 blocks in vitro TNA of anti-PA immune sera

Human Toxin Neutralizing Antisera Elicited by Anthrax Vaccine Also Augmented by FcR Interaction




AVR-801 Human Reference Standard Serum anti-PA IgG Concentration

Human PBMC Cytokine Inhibition Assay

Alternative to Murine Macrophage Assay

Whole blood




PharmAthene

MEDAREX

Human PBMC Cytokine Inhibition Assay


Toxin Dose Titration



Human PBMC Cytokine Inhibition Assay

Valortim Potency


TNF- α




Human PBMC Cytokine Inhibition Assay

Potency Augmented by FcR Interaction

IL-6 Production in PBMC's



Neutralization by ValortimTM



Role of FcR Engagement *in vivo*

- Lowest effective dose of Valortim to date, 1 mg/kg still provides serum levels in excess of amounts that F(ab)₂ fragment would work – will lower amounts work?
- Notable that protective polyclonal immune responses in 3 species (mouse, rabbit and human) show:
 - Strong dependence on FcR interaction for activity in TNA, AND
 - Undetectable blocking activity for attachment (FcR independent)
 - Suggests FcR interaction is an important attribute of natural protection
- Definitive experiments for this role will require:
 - In vivo experiments with F(ab)₂ fragments
 - Experiments in animals with defined FcR mutations
 - Mutants of Valortim with FcR binding mutations

Summary

Valortim has an optimum profile for an anthrax anti-toxin

- Human monoclonal antibody
- High affinity and specificity
- Potent neutralizing activity in vitro and in vivo
- Mechanism of action same as vaccine induced immunity and distinct from other Mabs
- In vivo efficacy at doses as low as 1 mg/kg

Acknowledgements

Valortim Product Development

MEDAREX



NIAID



PharmAthene

Valortim R&D Program

Medarex, Inc.

Laura Vitale

Tom O'Neill

Diann Blanset

Joel Goldstein

Xi-Tao Wang

Tibor Keler

Geoff Nichol

Dartmouth Medical School

Ron Taylor

Michael Fanger

USAMRIID

Gerry Andrews

Jeff Adamovics

Stephen Little



PharmAthene

MEDAREX